

**AC033**

Adult Cardiology & Clinical Cardiovascular Medicine • Poster Presentation

LONG-TERM OUTCOME OF PERIPARTUM CARDIOMYOPATHY AT THE UNIVERSITY COLLEGE HOSPITAL IBADAN

Anuoluwapo Makinde^{1,2,3,*}, Olanike Orimolade^{1,2,3}, Joshua Akinyemi⁴, Chidinma Ogah³, Ifunanya Nebo³, Abdulhammed Babatunde³, Omokorede Ademowo-Olusanya^{2,3,5}, Akinyemi Aje¹, Oladimeji Adebayo^{2,5}, Adeola Abdul-salam^{6,7}, Gbolahan Obajimi^{6,7}, Bukola Adesina^{6,7}, Christopher Aimakhu^{6,7}, Abiodun Adeoye^{1,2}, Adewole Adebisi^{1,2}, Olulola Oladapo^{1,2}, Okechukwu Ogah^{1,2,3}

Institutional affiliations 1Cardiology Unit, Department of Medicine, College of Medicine, University of Ibadan, Ibadan, Oyo State, Nigeria 2Cardiology Unit, Department of Medicine, University College Hospital, Ibadan, Oyo State, Nigeria 3Cardiomyopathies, Heart Failure and Atrial Fibrillation Research, Cardiology Unit, Department of Medicine, College of Medicine, University of Ibadan, Ibadan, Oyo State, Nigeria 4Department of Epidemiology and Medical Statistics, Faculty of Public Health, College of Medicine, University of Ibadan, Ibadan, Oyo State, Nigeria 5Institute of Cardiovascular Diseases, College of Medicine, University of Ibadan, Ibadan, Oyo State, Nigeria 6Department of Obstetrics & Gynaecology, Faculty of Clinical Sciences, College of Medicine, University of Ibadan, Ibadan, Nigeria. 7Department of Obstetrics & Gynaecology, University College Hospital, Ibadan, Oyo State, Nigeria * Presenter Correspondence: Professor Okechukwu S Ogah, Cardiology Unit, Department of Medicine, University College Hospital Ibadan, PMB 5116, Ibadan, Nigeria

Stand: POSTER #AC033 | **Venue:** Poster Exhibition Hall & Gallery - Eko Convention Centre | **Online Ref:** <https://nigeriancardiacsociety.com/abstracts/AC033>

Background Peripartum cardiomyopathy (PPCM) is an important cause of maternal mortality and morbidity. There is limited data on outcomes and prognostic markers in patients receiving contemporary evidence-based therapy, particularly in southern Nigeria. The objective is to report the outcomes of PPCM in our setting. **Methods** We conducted a single-centre cohort study of 74 PPCM patients managed at the University College Hospital, Ibadan. The women were followed for a median of 16 months (range 1- 99 months). The primary endpoint was all-cause mortality. **Results** Seventy-four (74) patients with PPCM were seen between January 1, 2009, and December 31, 2022. The mean age of the women was 30.5 ± 6.9 years (range: 18-46 years). The mean BMI was 23.9 ± 3.8 Kg/m² while the mean gestational age was 38.9 ± 2.4 weeks. Eighteen (24.3%) women were readmitted at least once, while 21 (28.4%) died during the follow-up period. The presence of these factors was significantly associated with higher rates of all-cause mortality: lower body mass index (p=0.017), lower systolic and diastolic blood pressures (p=0.003 and 0.020, respectively), and larger LV internal diameter in diastole (p=0.014). **Conclusions** This study identified factors associated with poorer outcomes in our cohort. The presence of pregnancy-induced hypertension appears to be associated with better outcomes

Keywords: Women, Pregnancy, Cardiomyopathy, Heart failure, Outcome