



AC057

Adult Cardiology & Clinical Cardiovascular Medicine • Poster Presentation

UNDERUTILIZATION OF STATINS AND OTHER LIPID LOWERING THERAPIES AMONG HIGH-RISK CARDIOVASCULAR PATIENTS IN NIGERIA: A NATIONAL CROSS-SECTIONAL STUDY

Olumide Akinniyi Akinyele¹, Akinyele Taofiq Akinlade², Tajudin Adesegun Adetunji³, Oladimeji Adebayo⁴, Stella Chinyelu Alagbe⁵, Sebastine Oseghae Oiwoh⁶, Yekeen Ayodele Ayoola⁷, Tolulope Taiwo Shogade⁸, Opeyemi Olalekan Oni⁹, Umar Faruk Abdullahi¹⁰, Francisca Olufunmilayo Inofomoh¹¹, Yakubu Lawal¹², Adenike Christianah Enikuomehin¹³, Osahon Otaigbe¹⁴, Oluwatosin Ruth Ilori¹⁵, Yusuf Idris Abiola¹⁶, Dapo Sunday Oyedepo¹⁷, Tijani Idris Ahmad Oseni¹⁸, Atinuke Oluwabukola Kalada Richard¹⁹, Joel Olufunminiyi Akande²⁰, Emmanuel Ovundah Nyeche²¹, Adebimpe Olukemi Yusuf²², Ahidiyu Anaryu Mamza²³, Oladimeji Adededeji Junaid²⁴, Bolade Folasade Dele-Ojo²⁵, Damilola Adeyinka Akinfaderin²⁶, Oluseyi Ademola Adejumo²⁷, Adeseye Abiodun Akintunde²⁸.

Stand: POSTER #AC057 | **Venue:** Poster Exhibition Hall & Gallery - Eko Convention Centre | **Online Ref:** <https://nigeriancardiacsociety.com/abstracts/AC057>

Background: Lipid lowering therapies (LLT), particularly statins, are recommended in established high cardiovascular risk patients. However, national data on LLT utilization remain limited despite growing cardiovascular disease burden. We evaluated prevalence, pattern, and determinants of statin use among high-risk cardiovascular patients in Nigeria.

Methods: We conducted a national cross-sectional study through the Nigeria Clinical Lipid Research (NiCLiR) Network across 15 secondary and tertiary healthcare facilities between August 2025 and May 2026. Adults (≥18 years) with established high-risk cardiovascular conditions, were consecutively recruited during routine outpatient visits. Standardized case report forms captured demographic, clinical, and treatment data. Multivariable logistic regression identified factors associated with statin use, reported as adjusted odds ratios (aORs) with 95% confidence intervals (CIs).

Results: Among 1,470 participants, (mean age was 57.3 ± 14.5 years; 46.2% male), 536 (36.6%) received statins. Among statin users, atorvastatin was the most prescribed statin (54.3%), followed by rosuvastatin (28.9%) and simvastatin (17.7%); 23.5% received high-intensity statins. Non-statin LLT combination (0.34%) was rare. Increasing age (aOR 1.013; 95% CI 1.005–1.021; p = 0.002), hypertension (aOR 0.286 for no vs. yes; 95% CI 0.191–0.429; p < 0.001), and family history of diabetes mellitus (aOR 0.417 for no vs. yes; 95% CI 0.298–0.582; p < 0.001) were independently associated with statin use.

Conclusion: LLT use among high-risk cardiovascular patients was low, with limited use of high-intensity statins. These findings highlight gaps in evidence-based lipid management and support interventions to improve risk-based prescribing, and guideline implementation

Keywords: Statin, Lipid lowering therapy, Cardiovascular disease, Cardiovascular risk, Guideline adherence, Nigeria