

**IC002****Interventional Cardiology & Catheterization** • Poster Presentation**CORRELATION BETWEEN INVASIVE AND NON-INVASIVE PULMONARY PRESSURE ASSESSMENT IN LAGOS STATE UNIVERSITY TEACHING HOSPITAL**

Oluwaseye M. Oladimeji¹, Alaba Oladimeji², Anthony G. Kweki³, Obawole M. Olalekan[?], Adeola O. Ajibare[?], Ramon K. Moronkola[?], Esther O. Ijaola[?], Oluwaseun O. Okunuga[?], Ajibola Y. Sanda[?], Onome A. Ehizojie[?], Oluronke T. Bamgboye[?], Kabirat I. Edun[?], Aishah O. Salaudeen[?], Timilehin E. Idowu[?], Alaba P. Adebola[?], Folashade A. Daniel[?], Michael O. Balogun[?], Tayo A. Addo[?], Oladipupo Olafiranye[?], Ijeoma Ekeruo[?], Oluwaseyi Bolorunduro[?], Obinnaya Emerole^{1?} ¹ Internal Medicine, Lagos State University Teaching Hospital, Lagos, NGA; ² Cardiology, Lagos State University Teaching Hospital, Lagos, NGA; ³ Internal Medicine/Cardiology/ICENI Fellow, Colchester Hospital, ESNEFT, Colchester, GBR; [?] Internal Medicine, Lagos State University Teaching Hospital, Lagos, NGA; [?] Internal Medicine, Lagos State University College of Medicine, Lagos, NGA; [?] Internal Medicine, Obafemi Awolowo University Teaching Hospitals Complex, Ile-Ife, NGA; [?] Internal Medicine, The University of Texas Southwestern Medical Center, Texas, USA; [?] Internal Medicine, Texas Cardiac Arrhythmia, Texas, USA; [?] Department of Cardiology, Mary Washington Healthcare, Woodbridge, USA; ^{1?} Internal Medicine, Georgia Heart Physicians Macon, Georgia, USA.

Stand: POSTER #IC002 | **Venue:** Poster Exhibition Hall & Gallery - Eko Convention Centre | **Online Ref:** <https://nigeriancardiacsociety.com/abstracts/IC002>

Background: Pulmonary hypertension (PH) is associated with substantial morbidity and mortality. Although right heart catheterization (RHC) remains the gold standard for diagnosing PH, it is invasive and often unavailable in resource-limited settings. Echocardiography is widely used for non-invasive assessment, but its diagnostic performance relative to RHC has not been adequately validated in this setting.

Methodology: In this prospective cross-sectional study, 102 patients with suspected PH were recruited. Participants underwent transthoracic echocardiography followed by RHC within one hour. PH was defined as a mean pulmonary arterial pressure >25 mmHg on RHC. Correlation and diagnostic accuracy analyses were performed.

Results: Seventy-one participants completed both investigations. PH was identified in 48 (67.6%) participants by echocardiography and 39 (54.9%) by RHC. Echocardiographic pulmonary artery systolic pressure demonstrated a moderate positive correlation with invasive measurements ($r = 0.635$, $p < 0.001$). Mean pulmonary artery pressure estimated using the Chemla and Syed formulae showed the strongest correlation with invasive measurements (both $r = 0.650$, $p < 0.001$), while pulmonary acceleration time-derived estimates demonstrated a weaker correlation ($r = 0.473$, $p < 0.001$). Echocardiographic PASP demonstrated a sensitivity of 86.9%, specificity of 52.0%, positive predictive value of 76.9%, and negative predictive value of 68.4%.

Conclusions: Echocardiographic estimation of pulmonary arterial pressure demonstrated moderate agreement with invasive measurements and high sensitivity for detecting PH. While echocardiography is a valuable screening tool, RHC remains essential for definitive diagnosis.

Keywords: Echocardiography, non-invasive assessment, pulmonary arterial pressure, pulmonary hypertension, right heart catheterization.